

Table 13: Categorisation of inhaled corticosteroids by dose – children*
(see also Figure 3)

ICS	Dose		
	Very low dose	Low dose	Medium dose#
Pressurised metered dose inhalers (pMDI) with spacer			
Beclometasone dipropionate			
Non-proprietary	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Clenil Modulite	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Qvar (extrafine) Qvar autohaler Qvar Easi-breathe	n/a	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day
Soprobec	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Ciclesonide			
Alvesco Aerosol inhaler	n/a	80 micrograms two puffs once a day	160 micrograms two puffs once a day
Fluticasone propionate			
Flixotide Evohaler	50 micrograms one puff twice a day	50 micrograms two puffs twice a day	125 micrograms two puffs twice a day
Dry powder inhalers (DPI)			
Budesonide			
Non-proprietary Easyhaler	n/a	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Pulmicort Turbohaler	100 micrograms one puff twice a day	100 micrograms two puffs twice a day 200 micrograms one puff twice a day	200 micrograms two puffs twice a day 400 micrograms one puff twice a day
Fluticasone propionate			
Flixotide Accuhaler	50 micrograms one puff twice a day	100 micrograms one puff twice a day	250 micrograms one puff twice a day
Mometasone			
Asmanex Twisthaler	n/a	200 micrograms one puff twice a day	n/a
Combination inhalers			
Budesonide with formoterol			
Symbicort Turbohaler	100/6 one puff twice a day	100/6 two puffs twice a day 200/6 one puff twice a day	n/a
Fluticasone propionate with salmeterol			
Combisal MDI	n/a	50/25 two puffs twice a day	n/a
Seretide Accuhaler	n/a	100/50 one puff twice a day	n/a
Seretide Evohaler	n/a	50/25 two puffs twice a day	n/a

* Different products and doses are licensed for different age groups and some are not licensed for use in children. Prior to prescribing, the relevant summary of product characteristics (SPC) should be checked (www.medicines.org.uk/emc).

Medium doses (shaded boxes) should only be used after referring the patient to specialist care.